

Masterarbeit  
MScPT 2017 ZHAW

Comparison of two commercially available devices to measure nitric oxide lung diffusing capacity in healthy, non-smoking adults, a randomized cross-over study.

Autor: Quintin de Groot

Betreuungsperson: Thomas Radtke, PhD<sup>1</sup>

Beteiligte Institutionen: Universität Zürich,

Institut für Epidemiologie, Biostatistik und Prävention

Rämistrasse 71, 8006 Zürich

Zürcher Hochschule für Angewandte Wissenschaften,

ZHAW, Institut für Physiotherapie, Technikumsstrasse 71,

8400 Winterthur

Datum: 17.05.2020

Anzahl Zeichen: 21853

<sup>1</sup>University of Zurich, Epidemiology, Biostatistics and Prevention Institute, Zurich, Switzerland

## Abstract

**Background:** Lung diffusion measurements are commonly used for lung function diagnostics. The usage of nitric oxide (NO), in addition to carbon monoxide (CO) during a lung diffusion measurement, could lead to a better understanding of the process of diffusion. **Research Question:** This study investigated whether the two on the market available instruments for DLNO measurement ('MasterScreenTM' (MS) OFT Pro (Jaeger, Switzerland); 'HypAir' (HA) (Medisoftware, Dinant, Belgium)), measure identical at the same time at the same place with the same protocol and with identical subjects. **Method:** This study followed a single-center randomized cross-over design carried out at the University Hospital of Zurich. 35 healthy subjects of mixed age and sexes were included and randomly allocated to perform slow spirometry, forced spirometry and lung diffusion measurements on either one of the devices. After completing all tasks, the participants performed the exact same maneuvers on the leftover device. The primary outcome was lung diffusion capacity for nitric oxide (DLNO). Secondary outcomes were lung diffusion capacity for carbon monoxide (DLCO), alveolar volume (VA) and breath hold time (BHT). Comparison of primary and secondary outcomes between both devices were calculated by using a linear mixed model. **Results:** The primary outcome DLNO showed a highly significant difference between the HA and the MS device. The mean difference was 24.1 mL/min/mmHg with a 95%CI of 21.8-26.4 mL/min/mmHg ( $p < 0.0001$ ). For DLCO the difference was not significant with a mean difference of -0.228 mL/min/mmHg (95%CI: -0.571 mL/min/mmHg and 0.116 mL/min/mmHg;  $p > 0.2$ ). The mean difference in alveolar volume between both devices was 0.484 L (95%CI: 0.447-0.512 L;  $p < 0.0001$ ). **Interpretation:** Although both devices show similar DLCO values, DLNO and VA showed systematic clinically relevant differences. There is no reasonable explanation to justify the difference.

**Clinical Trial Registration:** NCT04016597

## Abbreviation List

<b>BHT</b>	Breath hold time
<b>BMI</b>	Body-mass-index
<b>CO</b>	Carbon monoxide
<b>DLCO</b>	Lung diffusion capacity with carbon monoxide
<b>DLNO</b>	Lung diffusion capacity with nitric oxide
<b>HA</b>	Device: HypAir
<b>HE</b>	Helium
<b>ICC</b>	Intra class correlation
<b>MS</b>	Device: MasterScreen
<b>NO</b>	Nitric oxide
<b>SD</b>	Standard deviation
<b>SDws</b>	Within-subject standard deviation
<b>VA</b>	Alveolar volume
<b>95%CI</b>	95 percent confidence interval

**Introduction:**

Lung-diffusion measurement analyzes the ability to absorb gases the air we breathe into the bloodstream.<sup>1</sup> This is one of the standard procedures in lung function diagnostics along with the body plethysmography and spirometry and is used in a wide variety of pulmonary disorders like chronic obstructive pulmonary disease (COPD), asthma and cystic fibrosis.<sup>2-5</sup> In current lung diffusion measurement carbon monoxide (CO) is added to the inhaled air to measure the difference of particles after exhalation within the exhaled air (DLCO).<sup>6</sup>

In contrast to DLCO, the lung diffusion capacity measurement with nitric oxide (DLNO) also uses the gas nitric oxide (NO) in addition to CO. This technique allows a more differentiated view on various reasons of a hindered gas diffusion.<sup>7</sup> It could help to better understand problems during the gas diffusion. This procedure in combination with the existing DLCO method could lead to a better understanding of lung diseases and could become of great value in diagnostic procedures.

By now only a small number of references-values for the DLNO method have been published and most of these include a limited number of subjects.<sup>8-11</sup> In the study of Zavorsky et al. (2017) the authors pooled data from previously held studies by Aulani et al. (2010), van der Lee et al. (2006) and Zavorsky et al. (2008) to create a bigger and therefore more representative pool of reference-values.<sup>8,10-12</sup> This was done although these reference-values were created with different protocols and obtained by different devices. In the study done by Munkholm et al. (2018) the researchers compared their obtained data with every single set of reference-values pooled by Zavorsky.<sup>9</sup> They illustrate that every single set of reference-values shows different means of the DLNO values. The authors state that possible explanations for these differences might be different protocols, different equipment as well as different populations.<sup>9</sup>

So far, a protocol has been published to standardize the DLNO measurement to eliminate future methodological bias<sup>11</sup>. Still different equipment is used internationally to measure the diffusing capacity and therefore leave room for potential bias. There are two commercially available devices on the market: 'MasterScreenTM' (MS) OFT Pro (Jaeger, Switzerland) and

'HypAir' (HA) (Medisoftware, Dinant, Belgium). Both devices measure the NO concentration with an electrochemical cell.

A previously conducted pilot study, which aimed at a comparison of both devices reveals that the HA device constantly measures higher DLNO values (Radtke unpublished). Up to this point there has been no research published regarding a comparison of both devices, even though both are used within the field of research.

The aim of this study was to answer the question of whether the two on the market available instruments for DLNO measurement, measure identical at the same time at the same place with the same protocol and with identical subjects.

## **Method:**

### Design:

This study followed a single-center randomized cross-over design carried out at the University Hospital of Zurich. All included participants were invited to the research laboratory to perform spirometry as well as the diffusion capacity measurements. Subjects were informed about the project and asked for their consent to participate. After the randomization the participants performed the spirometry and the diffusion measurement on either one of the two devices. After completing all assessments on the first device the participants performed the exact same measurement on the second device. See section 'Protocol' for detailed information regarding the procedure.

### Participants:

This study included 35 healthy non-smoking adults of both sexes and varying ages ( $\geq 18$  years). Recruitment was done by personal invitation or email. The Epidemiology, Biostatistics and Prevention Institute at the University of Zurich, the physiotherapy master's program at the Zurich University of applied Sciences (ZHAW) were places for recruitment. Inclusion criteria were  $\geq 18$  years, Caucasian ethnicity. Exclusion criteria were smoker, individuals with known pulmonary or cardiovascular disease, acute respiratory symptoms, operations or radiation therapy of the chest, a body mass index (BMI)  $>30\text{kg/m}^2$  and pregnancy. The



subjects of the study live in Zurich or the surrounding area. Characteristics of the studied population are seen in Table 1.

**Table 1**

Patient characteristics at baseline

Variables	HA then MS	MS then HA	Complete
Subjects (n)	22 (63%)	13 (37%)	35
Sex			
male	11 (50%)	6 (46%)	17 (49%)
female	11 (50%)	7 (54%)	18 (51%)
Age (years)	40.8 $\pm$ 2.9 (22-62)	38.8 $\pm$ 5.2 (21-75)	40.0 $\pm$ 15.5 (21-75)
Height (cm)	175.5 $\pm$ 2.5 (159-200)	172.8 $\pm$ 2.7 (159-196)	174.5 $\pm$ 10.1 (159-200)
Weight (kg)	73.2 $\pm$ 2.4 (53-91)	63.5 $\pm$ 3.4 (47-91)	69,6 $\pm$ 12.4 (47-91)
BMI (kg*m <sup>2</sup> )	23.7 $\pm$ 0.6 (18.6-28.1)	21.1 $\pm$ 0.6 (17.1-23.7)	22,7 $\pm$ 2.8 (17.1-28.1)
RHR	71.6 $\pm$ 2.4 (50-95)	68.5 $\pm$ 3.0 (51-88)	70.5 $\pm$ 11.0 (50-95)
Spo2 (in %)	97.7 $\pm$ 0.2 (95-99)	97.3 $\pm$ 0.3 (94-99)	97.5 $\pm$ 1.0 (94-99)

Data are mean inclusive standard deviation and range. BMI, body mass index; RHR, resting heart ratio; Spo2, oxygen saturation; MS, 'MasterScreenTM' OFT Pro (Jaeger, Switzerland); HA, 'HypAir' (Medisoftware, Dinant, Belgium).

#### Randomization:

The allocation list was computer generated with the online randomization tool accessible at <http://www.randomizer.org>. Simple randomization has been used. The generated list contained numbers ranging from one to two. The number one implied the given subject started with the MS device and went on with the HA device. The number two indicated to start with the HA device and afterwards the MS device. The list was created by an independent person not involved in the study. Access to the list was restricted to two independent persons not involved in this study. Allocation concealment was ensured using central randomization, by

ad hoc request of the allocation sequence via phone. Blinding was not possible due to the protocol.

#### Ethics:

Since this study is of a methodological design focusing on a comparison of two devices and as it is not used for diagnostic purposes or treatment advice, it does not fall within the scope of the Human Research Act (HRA). Therefore, no authorization by the ethics committee was required. This has been examined and registered by the ethics committee of the canton of Zurich by means of a clarification of responsibilities (2019-02026).

#### Quality control:

The quality and reproducibility of the measurements were ensured by the following means. The devices were reviewed, and presets were made by technicians in advance of the study, to ensure that the devices were working properly. The software used for the HA device was 'Expair Version 1.32 05' for the MS device 'sentry Suite Version 3.0.2' was used. Each day the devices were manually calibrated using the three-flow method and a calibrated three liters syringe. Besides volume calibration, a gas calibration was performed using automated procedures for helium (He), carbon monoxide (CO), oxygen (O<sub>2</sub>), and methane (CH<sub>4</sub>). To ensure a detection of any major fluctuations in DLNO values during the course of the study, biological control measurements were performed. Therefore, the same subject performed every week all measurements on both devices, during the entire data collection period. This was done by two different subjects of both sexes (male, age: 26 years; female, age: 54 years).

#### Protocol:

In advance of the measurement, no intensive physical activities were allowed for at least 24 hours prior to testing. Coffee and meal consumption have been restricted three hours before undergoing the measurements. After welcoming the subject, the ad-hoc randomization has taken place by support of the statistical department. The aim of the study was explained to the subjects and the inclusion and exclusion criteria were checked. General information like bodyweight (nearest to 100g), standing height (nearest to 1 mm), sex, date of birth, resting heart rate (after a minimum of five minutes rest), oxygen saturation (in percent) as well as air

temperature (in degrees Celsius) and air-humidity (in percent) were noted. The measurements were taken at 430 meters above sea level. During the entire measurement, the subjects were asked to stay seated and could drink one cup of water in between the measurements. According to the randomization process, either one of the devices was used first. In the beginning, a slow spirometry was performed, and it was repeated until three reproducible measurements were collected. Afterwards, the forced spirometry was performed repeatedly until three reproducible measurements were gathered. Both spirometries were performed according to standardized protocols.<sup>13</sup> Following the forced spirometry, the lung diffusion capacity measurement was performed. Like the spirometry, standardized protocols were used for DLNO measurement and the maneuver was in the similar way repeatedly performed until three reproducible measurements (all DLNO scores within a range of 17 mL/min/mmHg), were collected.<sup>14</sup> In between the separate measurements a five-minute break as wash-out period was held. There were no changes to methods after trial commencement.

#### Outcomes:

The primary outcome of the measurements is DLNO (in mL/min/mmHg). Secondary outcomes are: DLCO (mL/min/mmHg), alveolar volume (VA; in liters) and breath-hold time (BHT; in seconds). Outcomes are obtained by stated DLNO devices. General information such as height, age and body weight were noted. Further values, for example BMI (body-mass-index) were calculated using patient characteristics.

#### Statistics:

Since there is no data available for of a power calculation, an a priori pilot study was used as a basis. From the values of the pilot study the ICC (Inter-Class-Correlation) for the DLNO values could be calculated as 0.96 (95% confidence interval (95%CI) 0.0.85-1). Because in the pilot study all participants were familiar with the procedure and that other external factors had an influence on the result, a more conservative approach was chosen for this study. For a desired ICC of 0.85 (95%CI 0.75-0.95) 31 participants were required (ICCest Calculation – Calculated with nQuery Advisor 7.0). Due to possible drop-out, 35 participants were recruited.

Comparison of primary and secondary outcomes between both devices will be calculated by using a linear mixed model (DLNO as primary endpoint) adjusting for device (Jaeger vs


Medisoft, coded as 0, 1) and period (machine with which the participant started) as fixed effects AND random intercept for each subject.

The SPSS version 23 (IBM Corp. Armonk, NY, USA) and R Version 3.6.3 (R-project, Vienna, Austria) are used for statistical evaluations. Three reproducible measurements per instrument (MS/HA) are at minimum required for calculation. The ICC 3.1 calculation (two way-mixed model) was used with its associated 95% confidence interval. Precision of DLNO values were be quantified by the within-subject standard deviations (SDws = root mean square error) calculated by the root-mean-square (RMS) method and the coefficient of variation.<sup>15</sup> Reproducibility was calculated with  $1.96 \times 1.96 \times \sqrt{2} \times \text{SDws}$  (95% confidence interval). Intra-device reproducibility was calculated as  $1.96 \times \sqrt{2} \times \text{SDws}$  (95% level of confidence).<sup>16</sup> Intraclass correlation coefficients (ICC's) and their 95% CI's were calculated for the primary outcome using a two-way mixed model [consistency, single measurement, (ICC, 3.1)].<sup>17</sup> Interpretation of ICC's is based on Cicchetti.<sup>18</sup>

## Results:

All participants completed all assessments and no participant was excluded from this study. No adverse events occurred during the entire data collection period. All participants were recruited and measured between October 2019 and January 2020. Trial was ended after completion of all measurements. No changes to outcomes after trial commencement were taken. See figure 1 detailed information about participant follow-up.

Figure 1  
Flow diagram




Flow diagram showing detailed information about patient follow-up. n, number of participants

Both sets of equipment passed all the tests performed. 22 Subjects were allocated to begin with the MA device, 13 allocated to begin with HA device. Subjects showed a mean resting heart rate of 70.5 (Standard deviation [SD]: 11.02; range: 50-95) and a peripheral capillary oxygen saturation of 97.7 (SD: 1.01; range: 94-99). Mean laboratory environmental conditions were: Temperature 23.7 °Celsius (SD: 1.66; range 21-27); Humidity 36.7 (SD: 8.96; range: 22.5-51). Allocation specific details are presented in Table 1.

The primary outcome DLNO difference between the HA and the MS device showed a highly significant difference. The mean difference was 24.1 mL/min/mmHg with a 95%CI of 21.8-26.4 mL/min/mmHg ( $p < 0.0001$ ). In figure 2 all DLNO results for each subject are plotted in ranked order.

Figure 2:  
Ranked DLNO results




All DLNO measurements per subject id in ranked order for DLNO. DLNO, lung diffusion capacity measurement with nitric oxide; Jaeger, MasterScreen™ (MS) device; Medisoft, HypAir (HA) device.

In contrast to DLNO, the DLCO measurement show no significant difference with a mean difference of -0.228 mL/min/mmHg and a 95% confidence interval between -0.571 and 0.116 mL/min/mmHg ( $p > 0.2$ ). The mean difference in alveolar volume between both devices was

0.484 L (95%CI: 0.447-0.512 L;  $p < 0.0001$ ). The average Breath-hold-time on the HA device was  $5.93 \pm 0.37$ , on the MS device the BHT was  $6.29 \pm 0.34$ . Figure 3 shows mean values on both devices for DLNO, DLCO and VA. Mean values on both devices are linear connected.

Figure 3

Linear connected means



Linear connected means between both devices. DLNO, lung diffusion capacity measurement with nitric oxide; DLCO, lung diffusion capacity measurement with carbon monoxide; VA, alveolar volume; Jaeger, MasterScreen™ (MS) device; Medisoft, HypAir (HA) device.

Intra-device variability characteristics are shown in Table 2. All Subjects fulfilled the recommended intrasession repeatability criteria for DLNO and DLCO (17.0 and 3.2 mL/min/mmHg).<sup>12,14</sup> The reliability analysis showed an ICC for the DLNO value of 0.979 with a 95% confidence interval of 0.959-0.990.

Table 2

Intra-session variability

Variables	MasterScreenTM (MS)			HypAir (HA)		
	Measure- ment error <sup>a</sup>	Repeatability <sup>b</sup>	CV (%)	Measure- ment error <sup>a</sup>	Repeatability <sup>b</sup>	CV (%)
DLNO mL/min/mmHg	6.80	18.84	4.9 9	6.26	17,35	3,74
DLCO mL/min/mmHg	1.17	3.23	3.8 2	1.05	2.92	3.41
VA (L)	0.11	0.30	1.7 8	0.11	0.30	1.78

Data were measured at a target breath-hold time of 5 s. CV, coefficient of variation; DLNO, pulmonary diffusing capacity for nitric oxide; DLCO, pulmonary diffusing capacity for carbon monoxide; VA, alveolar volume.

<sup>a</sup> The measurement error (or within-subject standard deviation, SDws) was calculated by the root-mean-square (RMS) method.

<sup>b</sup> Repeatability of gas diffusing measurements was calculated as the SDws from three single-breath tests for each study visit separately and multiplied by 2.77 (95% level of confidence).

During the time of data collection for biomonitoring, both selected subjects completed each eleven biomonitoring assessments. Both subjects showed only little variation within their scores during the entire assessment period. Person one showed on the HA device a coefficient of variation of 3.17 mL/min/mmHg and a mean deviation of 6.13 mL/min/mmHg for DLNO. On the MS device a coefficient of variation of 3.07 mL/min/mmHg and a mean deviation of 4.56 mL/min/mmHg. Person two showed a coefficient of variation of 2.72 mL/min/mmHg and a mean deviation of 2.6 mL/min/mmHg for DLNO. On the MS device person two showed a coefficient of variation of 3.06 mL/min/mmHg and a mean deviation of 2.64 mL/min/mmHg.



## Discussion:

The aim of this study was to compare two commercially available devices to measure DLNO.

This study shows that there is a statistically significant difference between both devices if comparing for DLNO. The within session repeatability for DLNO is about 17 mL/min/mmHg.<sup>14,19</sup> This study shows a higher mean difference (24.1 mL/min/mmHg; 95%CI: 20.5-27.6). It indicates that this difference is not only statistical but also clinically meaningful. In contrast to DLNO, DLCO show no statistically significant difference. The repeatability of DLCO is 2 -2.5 mL/min/mmHg.<sup>20</sup> Therefore, DLCO shows no clinical meaningful and no statistically significant difference between both devices. The coefficient of repeatability for VA for healthy subjects is 144 ml, consequently VA shows statistical significant and clinical meaningful differences.<sup>21</sup>

It is unclear how this systematic difference can be explained besides measurement differences. Both devices were running on the latest software update and have been controlled by technicians prior to this study, to ensure both are working properly. Both devices use similar electrochemical cells for the DLNO measurement. All instructions on both devices have been followed and the entire measurement procedure was held according to the latest protocol published by Zavorsky (2017).<sup>12</sup> A possible learning effect has been avoided by the cross-over study design. A possible carry over effect has been avoided by strict adherence to the wash-out period in between every single DLNO measurement and in between switching devices. Biomonitoring has been done during the entire data-collection period to detect any major fluctuations and have shown no major fluctuations whether in DLNO, DLCO or VA. Since there is no gold standard to compare these results with, these results cannot indicate whether one of both studied devices measures 'correctly'. Therefore, no right or wrong can be stated by this study.

In a previous study by Zavorsky and Murias (2007) the intra-session variability of DLNO and DLCO was analyzed for 31 healthy subjects using the HA device.<sup>19</sup> Their observed intra-session variability values for DLNO are comparable to the ones obtained in this study (Repeatability: 19.5 versus 17.35 mL/min/mmHg). There is no data available for intra-session variability for

the MS device for healthy subjects. However, in our study, both devices show overall comparable intra-session variabilities.

The difference in calculated BHT can be explained by different software interfaces and slightly different instructions. Studies have shown that shorter breath hold times can overestimate the diffusion capacity in healthy subjects.<sup>22,23</sup> The HA devices scored lower BHT values and higher DLNO values. The study of Dressel et al. (2008) showed no statistically significant differences in DLNO if comparing a BHT of four to six seconds.<sup>22</sup> A comparison between four and eight seconds showed a statistically significant difference with an absolute mean difference of 8,63 mL/min/mmHg.<sup>22</sup> In this study the mean difference in BHT was 0.36 seconds and the mean difference in DLNO was 24.1 mL/min/mmHg. Therefore, the difference in BHT could only explain a marginal amount of the presented difference.

This study used different gas canisters with a different amount of helium percentage (10%HE for the MS device; 14%HE for the HA device). The usage of a mixture containing 14% Helium was recommended by the 'MediSoft' company during the purchase of the device. Although it is recommended to use 10% HE there are also studies published for DLNO reference values using a mixture with 14% HE.<sup>10</sup> Helium is used as tracer gas and has no interference with the measurement of DLNO. For the calculation of the alveolar volume a helium dilution technique was used on both devices. The dilution factor for HE was adjusted according to the mixture used for both devices.

Up to today only a small number of reference values for the DLNO method have been published, most of these include a limited number of subjects.<sup>8-11</sup> Most of these studies were performed in a single laboratory and used only a single device. To increase the amount of reference values, studies with larger samples sizes would be necessary. This would exceed the possibilities for a single laboratory. Therefore, multicentric studies must be performed to be able to test many subjects. Subsequently established data could be pooled to increase reference values. This study shows that it is not recommended to pool data collected by both devices (MS and HA) to create reference values as it is done before.<sup>12</sup> These would not represent true reference values for any given population.

In addition to these findings, it is at this point of time unclear if the same device presents similar values given the same circumstances for the same person at two different locations. This question must be answered in advance of pooling data from either one of the devices from different locations. These findings show that pooling data for reference values is currently not accurate. Further research and development must be conducted to allow pooling data from different locations.

One limitation of this study is the missing block randomization. This allowed the allocation to exceed a close to 50-50 percent distribution. In this study 63 % were appointed to start with the HA device. However, characteristics are well balanced along both sequence groups.

In conclusion, this study shows that the two on the market available devices for DLNO measurement ('MasterScreenTM' OFT Pro [Jaeger, Switzerland] and 'HypAir' [Medisoftware, Dinant, Belgium]) present within the same individuals systematic statistical differences for the lung diffusion capacity measurement with nitric oxide. There was no clinical meaningful or statistical significant difference found for DLCO measurement at the same time. There is no reasonable explanation to justify the total amount of detected difference between both devices.

### **Acknowledgments:**

This study was funded by the Epidemiology, Biostatistics and Prevention Institute at the University of Zurich in collaboration with the Zurich University of Applied Sciences (ZHAW). All equipment used in this study was bought independently. No third parties were involved in funding this study. The Author 'de Groot, Quintin' is the guarantor of the content of the manuscript, including the data and analysis. 'Radtko Thomas' designed the study and contributed to data analysis and interpretation. 'Sarah Haile' contributed to data analysis.

### **References:**

1. Artmann F, Hader C, Rühle K, Rasche K. Die Diffusionskapazität in der täglichen Praxis. *Atemwegs- Lungenkrankh.* 2009;35:10-17. doi:10.5414/ATP35010

2. Barisione G, Brusasco C, Garlaschi A, Baroffio M, Brusasco V. Lung diffusing capacity for nitric oxide as a marker of fibrotic changes in idiopathic interstitial pneumonias. *J Appl Physiol Bethesda Md* 1985. 2016;120(9):1029-1038. doi:10.1152/japplphysiol.00964.2015
3. Chemery L, Fekete K, Guillot S, et al. Capacité de transfert du monoxyde de carbone (TLCO) et saturation artérielle à l'exercice chez les patients atteints de mucoviscidose. *Arch Pédiatrie*. 2004;11(9):1060-1066. doi:10.1016/j.arcped.2004.04.019
4. Espiritu JD, Ruppel G, Shrestha Y, Kleinhenz ME. The diffusing capacity in adult cystic fibrosis. *Respir Med*. 2003;97(6):606-611. doi:10.1053/rmed.2003.1487
5. Zavorsky GS, Wilson B, Harris JK, Kim DJ, Carli F, Mayo NE. Pulmonary diffusion and aerobic capacity: is there a relation? Does obesity matter? *Acta Physiol*. 2010;198(4):499-507. doi:10.1111/j.1748-1716.2009.02059.x
6. Graham BL, Brusasco V, Burgos F, et al. 2017 ERS/ATS standards for single-breath carbon monoxide uptake in the lung. *Eur Respir J*. 2017;49(1):1600016. doi:10.1183/13993003.00016-2016
7. Zavorsky GS, van der Lee I. Can the measurement of pulmonary diffusing capacity for nitric oxide replace the measurement of pulmonary diffusing capacity for carbon monoxide? *Respir Physiol Neurobiol*. 2017;241:9-16. doi:10.1016/j.resp.2016.11.008
8. Zavorsky GS, Cao J, Murias JM. Reference values of pulmonary diffusing capacity for nitric oxide in an adult population. *Nitric Oxide*. 2008;18(1):70-79. doi:10.1016/j.niox.2007.10.002
9. Munkholm M, Marott JL, Bjerre-Kristensen L, et al. Reference equations for pulmonary diffusing capacity of carbon monoxide and nitric oxide in adult Caucasians. *Eur Respir J*. 2018;52(1). doi:10.1183/13993003.00677-2015
10. Aguilaniu B, Maitre J, Glenet S, Gegout-Petit A, Guenard H. European reference equations for CO and NO lung transfer. *Eur Respir J*. 2008;31(5):1091-1097. doi:10.1183/09031936.00063207
11. van der Lee I, Zanen P, Grutters JC, Snijder RJ, van den Bosch JMM. Diffusing capacity for nitric oxide and carbon monoxide in patients with diffuse parenchymal lung disease and pulmonary arterial hypertension. *Chest*. 2006;129(2):378-383. doi:10.1378/chest.129.2.378
12. Zavorsky GS, Hsia CCW, Hughes JMB, et al. Standardisation and application of the single-breath determination of nitric oxide uptake in the lung. *Eur Respir J*. 2017;49(2). doi:10.1183/13993003.00962-2016
13. Miller MR. Standardisation of spirometry. *Eur Respir J*. 2005;26(2):319-338. doi:10.1183/09031936.05.00034805

14. Zavorsky GS. A small amount of inhaled nitric oxide does not increase lung diffusing capacity. *Eur Respir J*. 2006;27(6):1251-1257. doi:10.1183/09031936.06.00146805
15. Zavorsky GS. The rise in carboxyhemoglobin from repeated pulmonary diffusing capacity tests. *Respir Physiol Neurobiol*. 2013;186(1):103-108. doi:10.1016/j.resp.2013.01.001
16. Glüer C-C, Blake G, Lu Y, Blunt BA, Jergas M, Genant HK. Accurate assessment of precision errors: How to measure the reproducibility of bone densitometry techniques. *Osteoporos Int*. 1995;5(4):262-270. doi:10.1007/BF01774016
17. Bland JM, Altman DG. Measurement error. *BMJ*. 1996;312(7047):1654.
18. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol Assess*. 1994;6(4):284-290. doi:10.1037/1040-3590.6.4.284
19. Murias JM, Zavorsky GS. Short-term variability of nitric oxide diffusing capacity and its components. *Respir Physiol Neurobiol*. 2007;157(2-3):316-325. doi:10.1016/j.resp.2006.12.008
20. Punjabi NM, Shade D, Patel AM, Wise RA. Measurement Variability in Single-Breath Diffusing Capacity of the Lung. *Chest*. 2003;123(4):1082-1089. doi:10.1378/chest.123.4.1082
21. Prediletto R, Fornai E, Catapano G, Carli C. Assessment of the alveolar volume when sampling exhaled gas at different expired volumes in the single breath diffusion test. *BMC Pulm Med*. 2007;7(1):18. doi:10.1186/1471-2466-7-18
22. Dressel H, Filser L, Fischer R, et al. Lung diffusing capacity for nitric oxide and carbon monoxide: dependence on breath-hold time. *Chest*. 2008;133(5):1149-1154. doi:10.1378/chest.07-2388
23. Graham BL, Dosman JA, Cotton DJ. A Theoretical Analysis of the Single Breath Diffusing Capacity for Carbon Monoxide. *IEEE Trans Biomed Eng*. 1980;BME-27(4):221-227. doi:10.1109/TBME.1980.326726

## Appendix:

### 1: Autorenrichtlinien des gewählten Journals:


**CHEST®**

Official Publication of the American College of Chest Physicians

#### AUTHOR INFORMATION PACK

#### TABLE OF CONTENTS

•	<b>Description</b>	<b>p.1</b>
•	<b>Impact Factor</b>	<b>p.1</b>
•	<b>Editorial Board</b>	<b>p.1</b>
•	<b>Guide for Authors</b>	<b>p.2</b>



ISSN: 0012-3692

#### DESCRIPTION

*CHEST* is committed to advancing the care of patients served by multidisciplinary clinicians across pulmonary, critical care, and sleep medicine through the publication of clinical research relevant to today's challenges and reflecting advances on the horizon. To provide context in a fast-changing landscape, *CHEST* also incorporates review articles, offers commentaries, and fosters debate on emerging controversies. *CHEST* applies strict peer review standards to ensure the scientific rigor and publishes all content online within two weeks of acceptance.

#### IMPACT FACTOR

2018: 9.657 © Clarivate Analytics Journal Citation Reports 2019

#### EDITORIAL BOARD

##### EDITOR IN CHIEF

**Peter J. Mazzone, MD, MPH, FCCP**, Cleveland, OH, USA

##### ASSISTANT EDITOR

**Hilary Petersen, MPAS, PA-C**, Cleveland, OH, USA

##### DEPUTY EDITORS

**Susan Murin, MD, FCCP**, Davis, CA, USA

**Darcy Marciniuk, MD, FCCP**, Saskatoon, SK, Canada

**Christopher L. Carroll, MD, MS, FCCP**, Hartford, CT, USA

**All CHEST Editorial Board Members**

## GUIDE FOR AUTHORS

---

### GENERAL POLICIES & PROCEDURES

All manuscripts submitted to *CHEST* should be prepared in accordance with the instructions for authors, which adhere to the latest International Committee of Medical Journal Editors (ICMJE) [Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals](#).

### Animal Studies

For all animal studies, research must conform to the National Research Council guidelines, as well as local and state regulatory principles or requirements. Authorship

### Authorship

*CHEST* follows [ICMJE recommendations](#) for attributing authorship, which is based on the following 4 criteria: Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND Drafting the work or revising it critically for important intellectual content; AND Final approval of the version to be published; AND Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Individuals must meet ALL four criteria to be listed as authors. For more information on authorship, collaborators, group authorship, and responsibility for data, please visit the [National Library of Medicine's Fact Sheet on Authorship in Medline](#).

The final author lineup and order should be determined by all authors before submission and may not be changed without a written explanation and signed permission of all authors post submission. This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts: Before the accepted manuscript is published in an online issue: Requests to add or remove an author, or to rearrange the author names, must be sent to the Managing Editor from the corresponding author of the accepted manuscript and must include (a) the reason the name should be added or removed, or the authors' names rearranged and (b) written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal, or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author whose name is being added or removed. Requests that are not sent by the corresponding author will be forwarded by the Managing Editor to the corresponding author, who must follow the procedure as described above. Note that (a) the Managing Editor will inform the Journal Editors of any such requests and (b) publication of the accepted manuscript in an online issue is suspended until authorship has been agreed. After the accepted manuscript is published in an online issue: Any requests to add, delete, or rearrange author names in an article published in an online issue will follow the same policies as noted above and will result in a corrigendum.

### Communication

Communication related to submissions or submission inquiries should come from the corresponding author or principal investigator. Inquiries regarding manuscripts from non-authors, including inquiries from third-party medical writers and commercial medical writing companies, are strongly discouraged and may not receive a reply. conflict of interest

### Conflict of Interest

A conflict of interest is a financial or intellectual relationship or other set of circumstances that might affect, or reasonably be perceived by others to affect, an author's judgment, conduct, or manuscript. When in doubt, disclose. *CHEST* asks that authors report any potential conflicts in a three-year period prior to the date of submission and, if known, any upcoming conflicts. Categories to be reported include royalties or in-kind benefits (eg, travel, accommodations) from a commercial entity, shareholdings, speaker bureau activities, industry advisory committees, expert witness testimony, and litigation related to the subject of the manuscript.

Each author must report conflicts of interest in two places. Each author should provide a summary conflict of interest statement to be included on the **title page** of the manuscript. If an author has no conflicts of interest, a statement to this effect should be provided. Once the corresponding author has completed the submission process (including adding all co-authors and e-mail addresses into the appropriate fields), each author will be notified via e-mail that they need to sign in to their account



in ScholarOne Manuscripts and complete the combined electronic Authorship Confirmation/Conflict of Interest Disclosure form. Accepted manuscripts will not enter production or be scheduled for an issue until all forms for all authors are received.

It is important for conflict of interest to be reported in both places. The electronic form completed in ScholarOne Manuscripts serves as the official signed (electronically) documentation for the journal's records, whereas the summary statement included on the title page of the manuscripts is the statement accessible to the journal peer reviewers throughout the process and ultimately published in the Acknowledgments section of accepted articles in *CHEST*. Further, it is important to ensure that the information reported on the form and that disclosed on the title page match exactly. The Editorial Office will review this information and will contact the authors to remedy any discrepancies.

#### **Ethical Treatment of Patients/Subjects**

For all human research, authors must ensure that studies are in accordance with the amended Declaration of Helsinki. Authors should indicate in their manuscripts that they have obtained [informed consent](#) from patients for the procedure/treatment and for their medical data to be used in a study. IRB approval

#### **Institutional Review Board (IRB) Approval/Helsinki Declaration**

For any studies involving patients (including chart reviews), a statement of review and approval must be included. For example: "This study was conducted in accordance with the amended [Declaration of Helsinki](#). Local institutional review boards or independent ethics committees approved the protocol, and written informed consent was obtained from all patients."

The name of the committee and the approval number should follow this statement in the Methods section. If this is a multicenter study, the list may be provided in a separate Word document to be published as [Supplemental Material](#).

#### **Language (usage and editing services)**

Please write your text in good English (American or British usage accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's [WebShop](#) or [customer support site](#) for more information.

#### **Peer Review**

All submissions are subject to peer review. Authors are encouraged (and in the case of [Original Research](#) required) to provide the names of qualified reviewers who have had experience with the subject matter, but who are not affiliated with the same institution(s) as the author(s). *CHEST* will make the final selection of peer reviewers. Permissions

#### **Permissions**

Reuse of any previously copyrighted/published material, including material that appears on a website, within an article submitted to *CHEST* requires written permission from the copyright holder. Information on how to obtain permission for material published in *CHEST* is available via <https://journal.chestnet.org/content/permission>.

#### **Figures/Tables**

It is the author's responsibility to obtain written permission and, where necessary, pay any fees to the copyright holder for republication in *CHEST*. Obtain permission for all print, online, and licensed uses from the copyright holder (usually the publisher); Provide copies of the permission with the submission (attach it as supplemental material in the file upload area in ScholarOne Manuscripts); Acknowledge the source in the legend of the figure/table with a numbered reference; Provide the full citation in the reference list; and Ensure that any language requirements of the copyright holder have been met (eg, Reproduced with permission from XXX). Survey

#### **Survey Instruments/Questionnaires**

Patient-reported outcome (PRO) measures are increasingly used to assess different aspects of patients' health status, such as symptoms and health-related quality of life (HRQoL). For instance, results of HRQoL are frequently used as a primary outcome measure in [studies](#). Well-developed PRO measures are precise instruments that accurately assess patients' quality of life and by definition need to meet certain standards in terms of development and psychometric validation. Investigators should understand that "even small modifications can compromise the reliability and validity of these



instruments. Even modifications that may appear innocent, such as changing the format or layout of the instrument, changing the order of the items, or rewording the instructions, may alter the patients' responses. Therefore, it is important to administer only the exact version used in the validation."<sup>1</sup>

It is not acceptable to adapt, modify in any way, or translate these instruments into another language without the permission of the developer(s) of the instrument.<sup>1</sup> Lack of copyright does not imply that researchers are allowed to adapt, modify, or translate instruments.

Authors are responsible for obtaining permissions related to any survey instruments or tools relating to PRO used in their submission and for providing *CHEST* with a written copy of the permission to use (and modify or translate, if applicable) with the manuscript submission (attach it as supplemental material in the file upload area in ScholarOne Manuscripts). The Methods section of the paper should include a statement noting that permission has been obtained for use of the instrument or tool.<sup>2</sup>

References: Breugelmans MA. Dangers in using translated medical questionnaires: the importance of conceptual equivalence across languages and cultures in patient-reported outcome measures. *Chest*. 2009;136(4):1175-1177. Juniper EF. Medical questionnaires are copyrighted to ensure that validity is maintained. *Chest*. 2009;136(4):951-952. Consent

### **Preliminary Reporting of Data/Embargo**

*CHEST* does not consider the reporting of raw data or results, as required by funding bodies such as government institutions or commercial entities, to constitute prior publication. However, on acceptance for publication in *CHEST* journal, the article content is embargoed from media coverage and any media coverage should be coordinated through *CHEST*. More information is available in the [Media/Embargo Policy](#).

You may be invited by other organizations or companies to upload or share your submitted and/or accepted work. These companies are acting in violation of copyright law, and their activities may obscure the scientific record. Information on article rights is available at <https://www.elsevier.com/authors/journal-authors/submit-your-paper/sharing-and-promoting-your-article>. If you have questions about any requests received, please contact us at [editor@chestnet.org](mailto:editor@chestnet.org).

### **Privacy and Informed Consent**

Authors must omit from their text, tables, figures, and supplemental data any identifying details regarding patients and study participants, including names, initials, date of birth, Social Security numbers, dates, or medical record numbers (even when patient consent has been obtained). Authors must obtain written informed permission from the patient, guardian, or next of kin when individual cases are presented. Copies of the permission must be provided to *CHEST* prior to publication. If the patient has died or is otherwise unavailable, then permission must be sought from the next of kin.

Written patient permission to publish is required for all case-based sections of *CHEST*, including [Novel Reports](#), [Chest Imaging and Pathology for Clinicians](#), [CHEST Pearls](#), and [Ultrasound Corner](#). [Download the CHEST permission form here](#).

All authors are responsible for ensuring the submission complies with the [Health Insurance Portability and Accountability Act](#) or national equivalent.

### **Scientific Misconduct**

When *CHEST* has concerns, or receives allegations, of scientific misconduct, *CHEST* reserves the right to proceed according to the procedures described below and to the guidelines issued by the [Office of Research Integrity](#). *CHEST* recognizes its responsibility to appropriately address concerns and allegations of misconduct. Examples of misconduct include falsification of data, plagiarism (both plagiarism of others and self-plagiarism), improper designations of authorship, duplicate publication, misappropriation of others' research, failure to disclose conflict(s) of interest, and failure to comply with applicable legislative or regulatory requirements. Misconduct also includes failure to comply with any rules, policies, or procedures implemented by *CHEST* and other behaviors specified in the Office of Research Integrity guidelines.

*CHEST* follows the policies and recommended procedures outlined by the [Committee on Publication Ethics](#) in its approach to identifying and handling misconduct.

### Supplement Issue Proposals

Although *CHEST* will consider supplements sponsored by third parties for publication, it will publish only those supplements that advance the field or provide information that will significantly impact patient care in a novel way. **Proposal:** A complete draft table of contents, inclusive of titles, proposed authors, article lengths, and a brief description of what will be covered should be submitted to *CHEST* prior to the development of any further materials. Funding sources should also be disclosed. The material covered should have a broad interest to one or more constituents served by *CHEST* journal and *CHEST* (eg, pulmonologists, critical care physicians, cardiovascular or thoracic surgeons, and sleep physicians). The Editor in Chief will make a preliminary determination as to whether the proposal is of interest to *CHEST*. Final manuscripts will be submitted for peer review, and no guarantee of acceptance can be made. **Funding:** Supplements must have a commitment of funding, ideally from a nongovernmental organization, philanthropic foundation, or government-funded health-care body. The supporting organization shall not in any way dictate or impact the editorial content of the supplement. No title or article shall have the appearance of a conflict of interest, paid advertisement, or proprietary study. The Editor in Chief will make such determinations. **Draft Manuscripts:** Manuscripts should be written by the named authors. Any editorial assistance and/or writing support should be noted in the **acknowledgments** of each article, as should the source of funding for this assistance. Typically, one or more of the organizers of the supplement will provide a preliminary review of all the papers in a supplement for suitability of content, initial quality control, and adherence to agreed-on format. The format will be a coordinated effort of the supplement organizers and *CHEST*, who will work with authors before papers are formally submitted to the journal. Once the organizers have met their own standards for submission, they will provide the *CHEST* Editorial Office with a list of manuscripts, authors, and contact information for a Corresponding Author for each manuscript.

**Peer Review:** A designated supplement material receipt date will be set by the *CHEST* Editorial Office. All manuscripts and materials must reach the Editorial Office by that date. *CHEST* will contact all corresponding authors with instructions on finalizing and uploading manuscripts into the ScholarOne Manuscripts system. All *CHEST* requirements for authors also apply to authors of supplement papers. *CHEST* will send out all papers to external reviewers for evaluation. Authors will be responsible for making the requested changes. Decisions about publication will be made by the appropriate subspecialty Associate Editor and the Editor in Chief. **Editing:** *CHEST* will copyedit all articles for grammar and style. The corresponding author of each article will be responsible for review and approval of final page proofs. **Publication:** Publication date will be determined by *CHEST*. An estimated publication date will be set once *CHEST* offices have received all the supplement material. *CHEST* reserves the right to move up or delay publication. All supplements will appear online as a standalone issue of *CHEST*, available to all *CHEST* subscribers. **Reprints & Bulk Orders:** Single article **reprints, e-prints, and bulk orders** will be available on publication.

To submit a supplement proposal, contact *CHEST* at [editorialoffice@chestnet.org](mailto:editorialoffice@chestnet.org).

### Tobacco Policy

*CHEST* will not consider research and manuscripts that have been supported either directly or indirectly by tobacco companies.

### ACCEPTANCE AND PUBLICATION PROCESS

#### Proofing

After acceptance, one set of PDF page proofs will be emailed to the corresponding author. A link will be provided in the email so that the authors can download the files themselves. Elsevier provides authors with PDF proofs that can be annotated; for this you will need to download Adobe Reader version 9 (or higher) available free from <http://get.adobe.com/reader>. Instructions on how to annotate PDF files will accompany the proofs (also provided online). The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/reader/tech-specs.html>.

If you are not able to use the PDF annotations function, you may email a list of corrections (including replies to the Query Form) to Elsevier. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and either return via fax or scan and email the pages. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables, and figures. Significant changes to the article as accepted for publication will be considered at this stage only with permission from the Editor in Chief. *CHEST*



will do everything possible to get your article published quickly and accurately. It is important to ensure that all corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

### Copyright

Authors are welcome to submit manuscripts for exclusive publication in *CHEST*, provided they have not been published nor are under review elsewhere.

Upon acceptance of an article, authors will be asked to complete a journal publishing agreement (for more information on this and copyright, see <https://www.elsevier.com/copyright>). An email will be sent to the corresponding author confirming receipt of the manuscript together with the Journal Publishing Agreement form or a link to the online version of this agreement.

Authors may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <https://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article.

For open access articles: Upon acceptance of an article, authors will be asked to complete an Exclusive License Agreement (for more information see <https://www.elsevier.com/OAauthoragreement>). Permitted third-party reuse of open access articles is determined by the author's choice of user license (see <https://www.elsevier.com/openaccesslicenses>).

### Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. For more information, including guidelines regarding posting to local servers, see <https://www.elsevier.com/authors/journal-authors/submit-your-paper/sharing-and-promoting-your-article>.

Please be aware you may be invited by other organizations or companies to upload or share your submitted and/or accepted work. These companies are acting in violation of copyright law, and their activities may obscure the scientific record. If you have questions about any requests received, please contact us at [editor@chestnet.org](mailto:editor@chestnet.org).

### PubMed Central (NIH-Funded Work/Authors)

Studies (or authors) funded by the [National Institutes of Health](#) (NIH) must be deposited into PubMed Central. The publisher submits the final version of all articles (or authors) funded by the NIH to PubMed Central on the authors' behalf on publication. On submission of NIH-funded work, authors should and include the relevant grant numbers on the [title page](#).

### Open Access Options

*CHEST* offers an open access publishing option, with a publication fee, payable by authors or their research funder. When this option is exercised: The article will be made free to view on publication to all users of the *CHEST* websites with an icon denoting open access status; no login credentials or subscription will be necessary for access. *CHEST* will deposit the article into PubMed Central, where it will be indexed and publicly available. One of two Creative Commons Licenses will be assigned, depending on the open access option determined: **Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)**: Available for noncommercial purposes (\$3,000), this option lets others distribute and copy the article, and include it in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article. **Creative Commons Attribution License**. Authors may be required by the funding body to select this option (\$5,000). These articles will be assigned a [Creative Commons Attribution License](#) (CC BY) in place of traditional copyright. This allows all users of the article to copy, distribute, and/or adapt the work for commercial and noncommercial purposes, provided the user(s) attribute the source material by noting the citation of the original *CHEST* work. NOTE: This option is available only to those whose funding bodies mandate a CC BY license.

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design, in the collection, analysis, and interpretation of data; in the writing of the report, and in the decision to submit the article for publication. If the funding sources had no such involvement, then this should be stated.

*CHEST* also makes articles available to subscribers in developing countries and patient groups through the Elsevier access programs (<https://www.elsevier.com/access>). Online First

#### Online First

Most articles are posted online two to three weeks after acceptance. Online First articles will be updated as they move through the production process, but should not be considered final until they are published within an issue. These articles may have additional changes from Online First publication to the final, issue-assigned version. The article title, author names and affiliations, and abstract will be provided to PubMed for indexing. *CHEST* will not allow changes to the manuscript from the time of Online First publication until page proofs are received. Although Online First articles are indexed in PubMed, publication information will be updated as needed (ie, title change) at the time of final publication in a numbered issue of *CHEST*.

#### Online Only

The following sections are published online only: [Novel Reports](#); [Chest Imaging and Pathology for Clinicians](#); [CHEST Pearls](#); and [Ultrasound Corner](#). All online only content is indexed by PubMed, The Web of Science, and all search engines from which the majority of journal visits are derived. Online-only articles can be listed on a person's CV. The online journal is the journal of record and the primary archive. Manuscript Preparation

### GENERAL MANUSCRIPT PREPARATION

#### Submission

*CHEST* uses ScholarOne Manuscripts (<https://mc.manuscriptcentral.com/chest>) for manuscript submission and peer review. Submissions received by email or mail will not be considered. Technical assistance for ScholarOne Manuscripts is available by phone at +1-434-964-4100, and via online support (includes tutorials). Each submission is assigned a manuscript tracking number that will appear in the email that confirms your submission has been received. Please provide this tracking number on any correspondence regarding the manuscript.

When submitting to *CHEST*, authors will be asked to separately upload several distinct files through ScholarOne Manuscripts. The following list includes the types of files that may be required: Cover letter (either entered as text or uploaded to the Cover Letter area) Manuscript file (uploaded as Main Document), inclusive of **Title page Abbreviations list Abstract Text Acknowledgments References Tables Figure legends Figure files Supplemental material Permissions Patient consent**

#### Manuscript File

Title page

#### Title Page

The title page should be submitted as the first page of the main manuscript file and should include the following elements: **Word counts** for the text and abstract in the upper left-hand corner **Title** and short title/running head (of 50 characters or less) to be used in mobile formats **Author list** showing all names in the order and format that they are to appear on the publication. Also, include any middle initials and the highest degree obtained, as well as institutional affiliations. NOTE: Complete author information, including names, email addresses, and institutional affiliations must also be entered in ScholarOne Manuscripts to facilitate the collection of the required forms. **Corresponding author information** with full mailing address and email address (will appear on publication). Do not include phone or fax numbers on the title page. **Summary conflict of interest statements** for each author (or a statement indicating no conflicts exist for the specified author[s]) **Funding information** including any **NIH grant numbers** where applicable **Notation of prior abstract publication/presentation** including the name, date, and location of the relevant meeting

#### Abbreviation List

An alphabetical list of all abbreviations used in the paper, followed by their full definitions, should be provided on submission. Each abbreviation should be expanded at first mention in the text and noted parenthetically after expansion. Abbreviations should only be used for terms that appear more than three times in text. To aid readers, please use abbreviations sparingly.



### Abstract

For **Original Research** studies (clinical trials, interventional studies, cohort studies, case-control studies, epidemiologic assessments, surveys, systematic reviews, and meta-analyses), the abstract should consist of the following sections: Background Research Question Study Design and Methods Results Interpretation Clinical Trial Registration (registrar, website, and registration number), where applicable

For all other manuscript types requiring abstracts, CHEST requires a narrative (unstructured) abstract. More information is available in [Guidance for Specific Article Types](#).

### Text

#### Subheadings Within Articles

No more than 8 subheadings per article (in addition to headings such as Methods, Results, Discussion, and Interpretation). Each subheading can consist of only 5 words, including words such as a, an, the, and and.

### Acknowledgments

The acknowledgments section will vary slightly by article type. Possible elements include: **Guarantor** statement, naming one author who takes responsibility for (is the guarantor of) the content of the manuscript, including the data and analysis (**Original Research**) **Author contributions** should define the individual contributions each author made to the development of the manuscript and should include at minimum the three criteria required for **Authorship** as defined by *CHEST* (required for **Original Research**). If several authors made the same type of contributions, it is acceptable to combine them. An example author contribution line is: 'MLM had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis, including and especially any adverse effects. MLM, MT, NAW, DRG, VAD, and EG contributed substantially to the study design, data analysis and interpretation, and the writing of the manuscript.' **Financial/nonfinancial disclosures** should match those provided on the title page **Role of the sponsors** should detail what input or contributions, if any, were provided by the funding sources in the development of the research and manuscript **Other contributions** References

### References

Authors are responsible for the accuracy and completeness of citations. In text, references must be given as superscript numerals, numbered consecutively in the order in which they appear in the text. If the first (or only) mention of a reference appears in a Table, place the reference number after the Table call out in text. For example, if a reference is in Table 3 and has not been called out any earlier in the text, then the text call out should be, eg, "Table 3<sup>27</sup> ...". This will preserve numbering in citation management software. The full citations must be listed in numerical order at the end of the text. Each reference must contain, in order, the following: **Authors** (last name initials), listing all when there are up to six; first three followed by et al in the case of more than six authors

**Title** of article (sentence case, no quotation marks)

**Publication** source (italicized), when referring to a journal, the journal name should be abbreviated according to *Index Medicus*

**Year** of publication

**Volume** number

**Issue** number

**Page** numbers (inclusive)

### Journal Article

Sillen MJH, Speksnijder CM, Eterman R-MA, et al. Effects of neuromuscular electrical stimulation of muscles of ambulation in patients with chronic heart failure: a systematic review of the English-language literature. *Chest*. 2009;136(1):44-61.

Barker E, Haverson K, Stokes CR, Birchall M, Baily M. The larynx as an immunological organ: immunological architecture in the pig as a large animal model. *Clin Exp Immunol*. 2006;143(1):6-14.

### In-Press Journal Article

Annane D, Sebille V, Charpentier C, et al. Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock. *JAMA*. In press. <https://doi.org/10.1001/jama.288.7.862>

**Book**

Shields TW, LoCicero J III, Reed CE, Feins RH. *General Thoracic Surgery*. 7th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2009:200-232.

**Book Chapter**

Stone AC, Klinger JR. The right ventricle in pulmonary hypertension. In: Hill NS, Farber HW, eds. *Pulmonary Hypertension*. New York, NY: Humana Press; 2008:93-126.

**Abstract**

Garg N, Garg G, Christensen G, Singh A. Acute coronary syndrome caused by coronary artery mycotic aneurysm due to methicillin-resistant *Staphylococcus aureus* [abstract]. *Chest*. 2008;134(suppl):1001S.

For assistance in formatting other types of references, please refer to the American Medical Association Manual of Style.<sup>1</sup>

References: American Medical Association. *AMA Manual of Style: A Guide for Authors and Editors*. 10th ed. New York, NY: Oxford University Press; 2007.

**Open access**

Please visit our [Open Access page](#) for more information.

**Data References**

We encourage you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

**Tables**

Tables should be self-explanatory and should not duplicate text material. They must be numbered and cited in consecutive order in the text. Each must have a succinct title, column and row headings, and (where appropriate) a legend describing abbreviations and lettered footnotes at the bottom of the table. Tables should not contain any shading or special symbols and any special formatting (bold, italics) must be explained in the legend. Tables consisting of more than 10 columns are unacceptable and will not be published. Tables should be provided as word processing documents, not in a spreadsheet file format or as an image file. Tables may be added at the end of the main document file.

**Permissions** for any republished tables should be noted in the legend.

Tables used to describe or compare literature should include a column with the following information from the source publication: lead author last name, year of publication, and a numbered citation that corresponds to the full reference in the manuscript reference list.

**Figures**

Please make sure that artwork files are in an acceptable format (TIFF, JPEG, or EPS) and with the correct resolution. Color figures are published free of charge. For further information on figure preparation, please see <https://www.elsevier.com/artworkinstructions>.

Radiologic or other diagnostic examination figures or other diagnostic testing figures should have all patient-related numbering (including test date or medical record numbers) or wording removed prior to submission.

The journal encourages authors to supplement in-article microscopic images with corresponding high-resolution versions for use with the Virtual Microscope viewer. We also allow authors to enrich the online article by providing 3D radiological data in DICOM format. For more information on these Content Innovations, click [here](#).



### Figure Legends

All illustrations must be cited in consecutive numerical order within the text of the manuscript. A legend for each illustration should be provided on a separate page of the manuscript, not on the figure itself. Stains and magnifications for all photomicrographs should be included in the legend. Any image manipulation (eg, splicing) should be described in the legend. [Permissions](#) for any republished figures and any required [patient consent](#) lines for identifiable images also should be noted in the legend.

### Data visualization

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions [here](#) to find out about available data visualization options and how to include them with your article. Supplemental Material

### Supplemental Material/Appendices

Authors may submit supplemental material (ie, material that will be published only with the online version of the journal) if it enhances a study. The main text must stand alone, and the use of supplemental material should be judicious.

The same standards for ethics, copyright, permissions, and publication quality for the full-text article apply to all supplemental material. If any of the material included as supplemental material has been previously published, the authors are responsible for obtaining the required [permissions](#) and attributing the source material.

Appendices no longer appear in *CHEST* articles, but may be included as supplemental material, labeled e-Appendix. Lists of study participants and multicenter institutional review board data are examples of content that is appropriate for e-Appendices.

### Data visualization

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions [here](#) to find out about available data visualization options and how to include them with your article.

### Numbering

Each component of the supplemental material should be numbered and cited in consecutive order in the text of the article. Authors should not intersperse supplemental material consecutively with material for the print edition. The following convention should be used for labeling and numbering material: **e-Table:** number as e-Table 1, e-Table 2, etc

**e-Figure:** number as e-Figure 1, e-Figure 2, etc

**e-Appendix:** number as e-Appendix 1, e-Appendix 2, etc

**Audio:** number as Audio 1, Audio 2, etc

**Video:** number as Video 1, Video 2, etc (note, if shorter videos are combined into a single file, label each portion, eg, Video 1A, Video 1B, etc.)

Example: The distribution of missed bronchoscopy skills data points across centers and bronchoscopy milestones are depicted in e-Figure 1.

### Formats

The manuscript title, author list, and heading Supplemental Material should be included at the beginning of each file. The following formats can be uploaded as Online Content Only in ScholarOne Manuscripts: **Video:** Quicktime (.mov), Windows media (.wmv), Audio Video Interleave (.avi), animated GIF (.gif), .mpeg, and .mp4. All movie clips should be provided at the desired size and length (10 MB or 5 min maximum). Before submitting, authors should verify that clips are viewable in Quicktime or Windows Media Player. In addition, a brief text description should be provided in a word processing document explaining the video. Authors are encouraged to supply a still image of the video file for inclusion as reference in the print version of the article

**Audio:** .mp3, .wav, .au. In addition, a brief text description should be provided in a word processing document explaining the audio file.

**Tables:** Must be provided as Word files.

**Figures:** .tiff, .png, .jpeg, and .gif. One text document (in Microsoft Word) should be provided that contains brief captions for all figures.

**Text:** Microsoft Word (.doc, .docx), .rtf, and .txt files.

## References

References in supplemental material should be numbered consecutively beginning with 1; if a reference appears in both the main article and the supplemental material, it will likely have a different reference number. Supplemental material should be thought of distinctly in this regard.

## Style and Usage

*CHEST* follows the [AMA Manual of Style](#) (10th ed) in matters of editorial style and usage. All accepted manuscripts are subject to copyediting for conciseness, clarity, grammar, spelling, and *CHEST* style.

## Use of inclusive language

Inclusive language acknowledges diversity, conveys respect to all people, is sensitive to differences, and promotes equal opportunities. Articles should make no assumptions about the beliefs or commitments of any reader, should contain nothing which might imply that one individual is superior to another on the grounds of race, sex, culture or any other characteristic, and should use inclusive language throughout. Authors should ensure that writing is free from bias, for instance by using 'he or she', 'his/her' instead of 'he' or 'his', and by making use of job titles that are gender neutral (eg, "chairperson" instead of "chairman" and "flight attendant" instead of "stewardess"). Guidance

## GUIDANCE FOR SPECIFIC ARTICLE TYPES

In addition to following the general manuscript preparation instructions, authors should refer to the specific instructions for the type of article they are submitting.

**1 Section Consider Unsolicited (Y/N<sup>a</sup>) Abstract (wd max) Text<sup>b</sup> (wd max) References (max)**  
**Original Research** Y 300 3,200 50 **Guidelines and Consensus Statements** Y 300 4,000 150  
**Invited Content<sup>a</sup>** *CHEST Reviews* N 250 3,500 75 *How I Do It* N 250 3,000 50 *Point/Counterpoint*  
*Editorials* N None 1,200 12 *Editorials* N None 1,000 12 *Special Features* N 250 3,500 75 **Case Series**  
*Novel Reports* Y 150 750 20 *Chest Imaging & Pathology for Clinicians* Y None 1,600 10 *CHEST Pearls*  
Y None 1,600 10 *Ultrasound Corner* Y None 1,600 10 *Correspondence Letter to the Editor* Y None 400  
5 *Response to Letter to the Editor* N None 400 5 *Research Letter* Y None 1,000 10 *General Interest*  
*Commentary and Announcement* Y None 1,000 5

<sup>a</sup>These article types are invited. Authors with ideas for topics are encouraged to contact *CHEST* with proposals at [editorialoffice@chestnet.org](mailto:editorialoffice@chestnet.org). <sup>b</sup>Text word counts exclude abstract, references, figure legends, and tables. Original Research

## Original Research

**1 Article Element Requirements** Abstract length 300 words, structured format, include clinical trial information for randomized controlled trials Text length 3,200 words Reference count 75 references

## Format

A structured abstract should be provided. The abstract should be divided into the following sections: Background, Research Question, Study Design and Methods, Results, Interpretation, and Clinical Trial Registration Number (where applicable). The body of the text should be divided into the following sections: Introduction (not labeled), Methods, Results, Discussion, and Interpretation. Acknowledgements can follow (including author guarantor statement and contributions), then References. Finally, a Take Home Point pullout will be published. Please provide a sentence for the Study Question, Results, and Interpretation.

## Institutional Review Board (IRB) Approval

Most Original Research manuscripts must include a statement relating to institutional review board (or equivalent) approval in the "Methods" section. *CHEST* requires that authors include the committee name and approval number. In multicenter studies, the list of relevant committees and approval numbers may be included as an [e-Appendix](#). See more information on [IRB approval](#) here.

## Randomized Controlled Trials (RCTs)

*CHEST* defines a randomized controlled trial (RCT) as "any research study that prospectively assigns human participants or groups of humans to one or more health-related interventions to evaluate the effects on health outcomes." Authors preparing reports of RCTs for submission to *CHEST* should follow the [CONSORT \(Consolidated Standards of Reporting Trials\) checklist](#) and must include a CONSORT flowchart as Figure 1. Templates for the generation of CONSORT flowcharts are available [online](#).



In addition to following CONSORT, *CHEST* requires investigators to register their clinical trials in an approved public trials registry (see [Registration of Clinical Trials and Systematic Reviews](#) below). Purely observational studies (those in which the assignment of the medical intervention is not at the discretion of the investigator) do not require registration.

### **Systematic Reviews and Meta-analyses**

Authors preparing systematic reviews and meta-analyses for submission to *CHEST* should follow the [PRISMA \(Preferred Reporting Items for Systematic Reviews and Meta-analyses\) checklist](#) and must include a PRISMA flow diagram as Figure 1 on submission. *CHEST* strongly encourages registration of systematic reviews with the [PROSPERO registry](#) (see [Registration of Clinical Trials and Systematic Reviews](#) below). Additionally, authors are expected to address all items in the checklist in the writing of the manuscript. Those seeking additional guidance regarding the preparation of a systematic review can also consult the Cochrane Handbook for Systematic Reviews of Interventions at <http://www.cochrane.org/handbook> and the Institute of Medicine's Standards for Systematic Reviews available at <http://www.nationalacademies.org/hmd/Reports/2011/Finding-What-Works-in-Health-Care-Standards-for-Sy> of Trials

### **Registration of Clinical Trials and Systematic Reviews**

Authors of reports of clinical trials and systematic reviews should record their investigations in a viable registry (eg, [ClinicalTrials.gov](#), PROSPERO [<https://www.crd.york.ac.uk/prospERO/>]). Approved public trials registries are those that meet the criteria established by the [World Health Organization \(WHO\)](#). To register a trial, authors must submit the details directly to any one of the [WHO primary registries](#). *CHEST* reserves the right to reject papers if it deems the disclosure at the registry to be incomplete. An IRB statement is not a substitute for an approved clinical trial registration.

Authors should update their registrations to reflect any changes in outcomes, including primary and secondary end points, or protocols before participants are enrolled. The methods described in the published report must accord with those previously published in the study registration to avoid even the appearance of scientific misconduct. Furthermore, any changes to the original registration (eg, substituting a secondary outcome as the primary outcome) should be described in detail in the Methods section of the manuscript. Authors who modify their methods should post those changes on the online registry before submitting their manuscripts to *CHEST*. Studies

### **Surveys/Questionnaire-Based Studies**

Investigators who administer [surveys and questionnaires](#) as part of their study should obtain copyright permission if needed; no surveys should be adapted without the permission of the developer. Any unapproved changes in how PRO instruments are used or approved changes that have not been psychometrically studied and found to be reliable and valid will invalidate the results.

Authors of studies based on surveys or questionnaires should report on data that have been collected within two years of submission, including supporting reliability and validity data. All survey-based studies should describe the method used to achieve the response rate (eg, Dillman's tailored design method) and should provide a convincing rationale for why lower response rates provide important and generalizable information. Nonrespondents should be characterized well enough to allow for assessment of potential for nonresponse. Authors are encouraged to report outcome rates for most surveys using standardized definitions and metrics (eg, those proposed by the [American Association for Public Opinion Research](#)). This information must be detailed in the methods section.

### **Other Study Types**

The [Equator Network](#) provides checklists for other types of studies such as the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) statement. Checklists are also available for cohort, case-control, and cross-sectional studies, and authors are encouraged to follow these.

### **Confidence Interval**

For clinical studies, the primary outcome should be expressed as the difference between groups with a confidence interval (CI) around that difference provided in the Abstract and in the main article. In most cases, P values should not be presented without an accompanying effect estimate and CI. The CI is useful to readers because it indicates the precision of an estimated population value.

### Matching Language to Level of Evidence Guidelines

CHEST endorses the HEART Group Statement<sup>1</sup> calling for better matching language in original research to the evidence found in different study designs.<sup>2</sup> In short, in observational studies investigators should use descriptive statements such as “we observed a lower risk” rather than a more definitive statement such as “reduced the risk by” that are more appropriate to RCTs. Editors of Heart Group Journals. Statement on matching language to the type of evidence used in describing outcomes data. *J Am Coll Cardiol.* 2012;60(23):2420. Kohli P, Cannon CP. The importance of matching language to type of evidence: avoiding the pitfalls of reporting outcomes data. *Clin Cardiol.* 2012;35:714-717. Guidelines

### Guidelines and Consensus Statements

**1 Article Element Requirements** Executive summary Provided in bold text and including one to two paragraphs of introduction, followed by a summary of the data and a bulleted list of all recommendations and suggestions included in the document Abstract length 300 words, structured format Text length 4,000 words (may be negotiated with CHEST)

CHEST Guidelines are generated by CHEST under a well-defined development process. Committees work closely with the Guideline Oversight Committee, the Editor in Chief of CHEST, and relevant Associate Editors in developing guideline articles intended for submission to CHEST.

CHEST will work with other guideline-producing organizations where the possibility of mutual benefit exists. This includes guidelines and consensus statements where CHEST (the organization) has either agreed to participate in the development process, or has agreed to endorse the guideline or statement, or has been uninvolved in the development process.

Guideline-producing organizations that are not connected to a journal may submit their guideline for publication in CHEST. The submission will undergo peer review to include review by an internal methodologist and member of the CHEST Guideline Oversight Committee. For guidelines produced by organizations that publish in a subspecialty journal, a summary of the guideline publication with implementation tools may be published in CHEST with the goal of reaching our broad clinical audience. For these types of projects, authors should Contact the Editor in Chief of CHEST prior to submission. Recognize that a formal review of the submission will take place before a publication decision is made. Consider including implementation tools in the document or as a supplement. The Institute of Medicine's Standards for Developing Trustworthy Clinical Practice Guidelines should be consulted for guidance when using systematic reviews as the basis for guideline recommendations; these are available at <http://www.nationalacademies.org/hmd/Reports/2011/Clinical-Practice-Guidelines-We-Can-Trust.aspx>. Invited Content

### Invited Content

Reviews

### CHEST Reviews

**1 Article Element Requirements** Abstract length 250 (narrative) Text length 3,500 words Reference count 75 max

CHEST reviews in “sub-specialty” (eg, “Asthma”, or “COPD”) can include clinical, translational, ethics, education, and practice management topics. When a topic does not neatly fit in a sub-specialty section it may be included in the [Special Features](#) section (see below).

CHEST reviews are state-of-the art concise reviews on focused clinical, translational, ethics, education, and practice management topics. These should include a description of the importance of the topic and a summary of what is known about the topic with special attention to the most recent advances impacting practice. When relevant, authors should consider including sections on anticipated future advances, the authors' perspective on the topic, and summary tools that could assist with the application of the review in practice (eg, summary tables, algorithms). Topics in this section are developed and invited by the CHEST Associate Editors and Editor in Chief. Authors with suggestions for a topic are encouraged to contact CHEST at [editorialoffice@chestnet.org](mailto:editorialoffice@chestnet.org). How I Do

### How I Do It

**1 Article Element Requirements** Abstract length 250 words (unstructured) Text length 3,000 words Reference count 50 max



The How I Do It Section includes practical reviews of well-defined clinical questions with tools to assist with addressing the question when faced in practice. A relevant clinical question may have good evidence and guidelines available to support the approach, but implementation assistance is needed, or have weak evidence to support an approach but is a question with which clinicians struggle in practice.

Articles should be organized as follows: A well-defined clinical or procedure-related question. Case example. Review of relevant literature. Comment on nuances when applying to patient care. Review of relevant guidelines. Comment on nuances when applying to patient care. A summary table or algorithm whenever relevant. Summary of the approach to the question. Suggestions for multimedia adjuncts to the article are welcome.

Manuscript for the How I Do It section are invited by the subspecialty editorial teams.

Unsolicited contributions will not commonly be considered. Authors with ideas for topics should contact *CHEST* at editorialoffice@chestnet.org before preparing a manuscript.

### **Point/Counterpoint Editorials**

Point/Counterpoint editorials are submitted in two stages, each with distinct requirements: the Point and Counterpoint pieces have longer word limits, and the rebuttals are intended to be more succinct.

#### **Point/Counterpoint:**

**1 Article Element Requirements** Abstract length None Text length 1,200 words Reference count 12 references Figure/table limits 3 total tables and figures (not 3 of each)

#### **Rebuttals:**

**1 Article Element Requirements** Abstract length None Text length 500 words Reference count 7 references Figure/table limits 1 figure or table

Point/Counterpoint Editorials present a debate by content experts with different interpretations of the evidence supporting an approach to a topic. Authors on each side of the debate develop a rationale for their stance and then are given an opportunity to view and respond to the rationale provided counter to their stance.

Point/Counterpoint Editorials are invited by the editorial team. Authors with suggestions for a topic should contact *CHEST* at editorialoffice@chestnet.org prior to developing a manuscript.

### **Editorials**

**1 Article Element Requirements** Abstract length None Text length 1,000 words Reference count 12 max

Editorials are invited by the editorial team. They are meant to allow a content expert to discuss the findings of an original research article, sharing their perspective on how the publication advances the field, impacts practice, and highlights further research needs.

### **Special Features**

**1 Article Element Requirements** Abstract length 250 words, narrative format Text length 3,500 words Reference count 75 references

Special Features are invited reviews, commentaries, and other items of interest that do not fit well into other categories. NOTE: Systematic reviews should be submitted as [Original Research](#). *CHEST* will consider unsolicited Special Feature submissions, but authors must be aware that at any given time *CHEST* also has a long list of pending invited topics. Authors are encouraged to contact *CHEST* at editorialoffice@chestnet.org with a proposal on the topic prior to the writing or submission of any Special Feature articles.

### **Case Series**

Novel Reports

**Novel Reports (Online only)**

**1 Article Element Requirements** Abstract length 150 words, narrative format Text length 750 words, for a single case report; 1,600 words for multiple cases Reference count 20 references Format Introduction, Case Report(s), Discussion Other [Written patient permission is required for publication](#)

Case report submissions to *CHEST* should describe a new entity, mechanism, presentation, means of diagnosis, or treatment of a disease. All submissions to this section must be novel and/or unique. It is appropriate to submit a single case or multiple cases highlighting the same message. Studies with a research question that is addressed by a case series should be submitted as original research.

Case reports do not need institutional review board approval, but authors must preserve patient privacy and follow the [Health Insurance Portability and Accountability Act](#) or national equivalent rules in writing up the case. On acceptance, *CHEST* will require submission of [written patient permission](#) for publication. It is acceptable to submit case reports to *CHEST* that have been presented at meetings and congresses. This information should be disclosed on the title page and provided in the [references](#). Chest Imaging

**Chest Imaging and Pathology for Clinicians (Online Only)**

**1 Article Element Requirements** Abstract length None Text length 1,600 words (of which clinical, radiologic, and pathologic findings and discussion should be approximately 500 words each) Reference count 10 references Format Case Presentation (with distinct Clinical, Radiologic, and Pathologic Findings subsections); Q: What is the Diagnosis; A: Diagnosis; Discussion (with distinct Clinical, Radiologic, and Pathologic Discussion subsections); Bulleted list (3-4 lines at the most) of the take-home message from the case. Other [Written patient permission is required for publication](#)

Chest Imaging and Pathology for Clinicians is designed to aid readers in understanding the connection between clinical, radiographic, and pathologic features of a disease state. Each submission should include distinct clinical, radiologic, and pathologic sections within the case presentation and the discussion.

The format for submission to this section is as follows: **Title:** should include a short summary of the presenting feature, but not the diagnosis (ie, Dyspnea with slow-growing mass of the left hemithorax) **Case Presentation:** should include the following sections in sequence without the use of subheadings and without giving away the diagnosis: A clinical findings section should mention the relevant positives and negatives while avoiding detailed description of hospital course. The focus should be on the approach taken by the authors to make the diagnosis. Comments on the differential diagnosis and a table summarizing the clinical and radiologic features of the differential diagnoses are desirable. A radiologic findings section should briefly detail the plain chest radiograph (no corresponding figure need be submitted) and describe in detail the additional imaging studies performed, emphasizing findings that point to the diagnosis A pathologic findings section should describe these findings in detail and should focus on correlations with the radiologic findings. The pathology presented should confirm the diagnosis. Gross pathology or high-quality, low-power images that capture the radiologic and pathologic correlation are recommended. **What is the diagnosis?** Alternative questions may also be included (ie, What study should be conducted next?) in addition to the diagnosis question. **Diagnosis: XXX;** should also include the answer to any other questions posed **Discussion** should include the following sections in sequence with the use of subheadings **Clinical discussion** should illuminate how the clinical findings tie in with the diagnosis, addressing the typical and atypical case features. Authors are encouraged to highlight the clinical features that may alert the clinician to the diagnosis. In case of a rare disease, and brief description as well as diagnostic tests/criteria should be included. These may be tabulated. In the last paragraph, the outcomes of the case and the result of described intervention are useful. **Radiologic discussion** should highlight specific findings from chest radiographs and CT, PET, and MR scans. Authors are encouraged to highlight findings that exclude diagnosis and elaborate on the use of particular modalities. **Pathologic discussion,** should highlight pathologic patterns of lung involvement that correspond to patterns seen on chest imaging, and the pathologic differential diagnosis of the disease under discussion should be presented. Special staining techniques that may allow the diagnosis to be established should be addressed. **Conclusion:** a bulleted list (3-4 lines at the most) of the take home message from the case for clinicians is encouraged. **Image Quality Considerations**

**Sizing:** Images should be appropriately sized to minimize superfluous information—including, in particular, any surrounding structures outside the body. **Labeling/Figure legends:** Legends should include baseline information: slice thickness (in mm), orientation (axial, coronal), and reconstruction



algorithm (in the case of lung, either "smooth" or "edge enhanced" or their equivalent). For contrast-enhanced pulmonary artery studies, provide the rate, timing, volume and type of contrast as appropriate. Additional Imaging techniques: of particular interest is the addition of "movie" files (AVI or equivalent) when these augment image interpretation (eg, cardiac, aortic, or general vascular cases). The inclusion of other standard imaging formats, such as volumetrically rendered images and maximum (MIPS) and minimum (MinIPS) projection images, can be helpful. CHEST Pearls

### CHEST Pearls

**1 Article Element** Requirements Abstract length None Text length 1,600 words (of which case presentation should be under 300) Reference count Up to 10 references listed under a heading of "Suggested Readings." in chronological order; no citations in text. Format See below Other [Written patient permission is required for publication](#)

Manuscripts for this section are designed to present a case, pose a question, provide the answer, and summarize the main teaching points as Pearls.

**Title** should include a short summary of the presenting feature, but not the diagnosis. **History** should provide the recent clinical presentation with relevant past medical history, with enough information regarding relevant positives and negatives to allow construction of a reasonable differential diagnosis. **Physical Examination Findings** should give the patient's vital signs and other physical findings labeled according to organ system (eg, chest: bibasilar rales; cardiac: grade II/VI holosystolic murmur at the apex radiating to the axilla; abdomen: non-tender without organomegaly). **Diagnostic Studies**, should list all of the relevant normal and abnormal studies required to construct a reasonable differential diagnosis: hemogram, blood chemistry, urine studies, arterial blood gases, microbiology results, tissue biopsy studies, miscellaneous studies (ECG, esophageal motility studies, etc), radiographic studies, polysomnographic studies. Authors should place normal values in parentheses when referring to unusual test results or values that have different normal ranges between laboratories. **What is the diagnosis?** Additional questions may also be included (ie, What study should be conducted next?) in addition to the diagnosis question. Alternative questions may focus on management alone when a manuscript does not present a diagnostic question (eg, end-of-life management issues). **Diagnosis:** State the diagnosis and the answers to any additional questions posed in the preceding "What is the diagnosis?". Do not provide explanatory text here but just mention the answers.

**Discussion**, using the present tense, present a clear discussion of the clinical condition that flows clearly from one topic to another. Most manuscripts should cover sequentially the topics of epidemiology, pathophysiology/etiology, clinical manifestations, approach to diagnosis, treatment and outcomes. Exceptions, such as manuscripts on end-of-life decision-making, should retain a clearly organized sequence of topics. Do not refer to the present patient in the body of the general discussion but instead refer back to the present patient in the Clinical Course section. Avoid in the Discussion stating the findings or opinions of others (eg, Jones and Smith reported...); instead, authors should synthesize the literature and state their views on the topic. **Clinical Course**, should take the general discussion back to the specific patient presented, informing readers how the diagnosis was established, how the care was managed and what outcomes occurred. **Pearls**, 3 to 5 important teaching points extracted from the Discussion. Pearls should represent concise, specific and clinically useful information rather than general statements of fact. **Suggested readings**, should be listed in chronological order with the oldest first and include a mix of classic and recent journal or book citations. References to general medical or nursing textbooks should be avoided.

Figures are needed only for the case presentation. In discussing figures in the case report, simply refer to their presence when the findings are sufficiently obvious to challenge the reader. If the finding is subtle and difficult to detect, the abnormality can be described in the case report, but in describing the figure do not provide the diagnosis or the answer to the question you will pose in the manuscript. When not mentioned in the case report, the abnormality in the figure should be discussed in the body of the discussion on the following page when referring in general to the condition and in the section on clinical course when providing follow-up for the patient presented. Authors may consider including an algorithm describing an approach to the clinical presentation.

**Sample:** Brownback KR, Crosser MS, Simpson SQ. A 49-year-old man with chest pain and fever after returning from France. *Chest*. 2012;141(6):1618-1621. Ultrasound Corner

### Ultrasound Corner

**1 Article Element Requirements** Abstract length None Text length 1,600 words (of which case presentation should be up to 300 words, with the discussion being 900 words, including take-home points [ie, "Reverberations"]) Reference count 10; no references should appear before the Discussion Videos 2 or 3 video file sets (more than 1 video clip may be compiled for use in each video set),<sup>a</sup>: sets typically include 1) first step in diagnosis; 2) next step by ultrasonography or determination of diagnosis; 3) discussion video. Authors are responsible for creation and editing of videos, including addition of captioning and labeling.<sup>b</sup> Section editor will work with authors and CHEST to add voice-over narration of the discussion video on acceptance. Files names must be video1.XXX, video2.XXX, etc. and each Ultrasound Corner manuscript must have discussion video with the file name discussion.XXX (XXX is the file format). See past articles for the Discussion video format. Format 1) Introduction/case presentation + initial examination video set (do not describe the ultrasonogram in a manner that would provide the answer to your question in #2; do mention the part of the body from which the ultrasonogram was obtained); 2) One question + one answer and follow-up ultrasonography video set; 3) Discussion + discussion video; 4) 3-4 "reverberations" (ie, take-home points); 5) references; 6) captions for figures if included; 7) short description of each video Other [Written patient permission is required for publication; waivers may be considered on a case-by-case basis and must be approved by the Editor in Chief.](#)

<sup>a</sup>Video clips may be combined as needed.

<sup>b</sup>Authors should combine all needed video clips for each step into a single video file, using software such as Windows MovieMaker or Apple Final Cut Pro. For short ultrasound readings (eg, 2 or 3 seconds), authors should either loop the frames or copy the sequences several times so that viewers have a chance to absorb what they are seeing. Correspondence

### Correspondence

Letters

#### Letter to Editor, Response to Letter to Editor

**1 Article Element Requirements** Abstract length None Text length 400 words Reference count 5 references Other No [No supplemental material](#). One figure or table permitted.

Letters to the Editor are intended for the clarification and edification of articles published in *CHEST*. It is up to the discretion of the Editor in Chief whether any Correspondence is sent for external peer review and whether to accept any letter for publication.

#### Commenting on Recent Articles

All letters commenting on previous articles should strive to provide constructive and respectful comments of the original work. Letters should pertain to articles published within the preceding 6 weeks. Any correspondence discussing recent *CHEST* articles should include a short original title that does not duplicate the title of the article. Authors should include the full citation of the complete article in the reference list. For letters responding to articles published to the [Online First](#) section, *CHEST* will hold publication until the final version of the article is published in a numbered issue of *CHEST*. All accepted letters will be sent to the corresponding author of the original article with an invitation to submit a response for publication. Response

#### Response Letters

Authors are asked to submit all replies to letters on their work within 2 weeks of receiving the invitation. If they do not respond within this time frame, the original letter will be published without a response. Authors should never correspond directly with the authors of correspondence. The replying author should also include the full reference to their original work and should submit the same [conflict of interest](#) information relevant to the original work. *CHEST* reserves the right to update the conflict of interest line in this regard as needed. Research Letters

#### Research Letters

**1 Article Element Requirements** Abstract length None Text length 1,000 words Reference count 10 references Other No [No supplemental material](#); up to two figures and/or tables permitted.

Research Letters should be descriptions of focused research findings. The findings should be of high quality, be novel, or have potential clinical impact, but should not be advanced or large enough to warrant publication of a complete original research manuscript. Research Letters do not require an abstract. The text should include Introduction (not labeled), Methods, Results, and Discussion sections. They should follow the guidelines for [Manuscript Preparation and Submission Requirements](#). General Interest

**General Interest Commentary and Announcements**

**1 Article Element Requirements** Abstract length None Text length 1,000 words Reference count 5 references Other No [No supplemental material](#); one figure or table permitted.

*CHEST* will consider correspondence in the form of commentary of potential interest to readers, or that serves to announce matters of importance to the pulmonary, critical care, and sleep medicine community.

© Copyright 2018 Elsevier | <https://www.elsevier.com>



## 2. CONSORT Checklist:

## RESEARCH METHODS AND REPORTING

Table 1 | CONSORT checklist of information to include when reporting randomised crossover trials

Section/topic	Item No	Description	Page No*
Title†	1a	Identification as a randomised crossover trial in the title	0
Abstract†	1b	Specify a crossover design and report all information outlined in table 2	1
Introduction:			
Background‡	2a	Scientific background and explanation of rationale	2
Objectives‡	2b	Specific objectives or hypotheses	3
Methods:			
Trial design	3a	Rationale for a crossover design. Description of the design features including allocation ratio, especially the number and duration of periods, duration of washout period, and consideration of carry over effect	3
Change from protocol‡	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	6
Participant‡	4a	Eligibility criteria for participants	3-4
Settings and location‡	4b	Settings and locations where the data were collected	3
Intervention‡	5	The interventions with sufficient details to allow replication, including how and when they were actually administered	5-6
Outcomes‡	6a	Completely defined prespecified primary and secondary outcome measures, including how and when they were assessed	6
Changes to outcome‡	6b	Any changes to trial outcomes after the trial commenced, with reasons	8
Sample size‡	7a	How sample size was determined, accounting for within participant variability	6
Interim analyses and stopping guidelines‡	7b	When applicable, explanation of any interim analyses and stopping guidelines	6-7
Randomisation:			
Sequence generation‡	8a	Method used to generate the random allocation sequence	4
Sequence generation‡	8b	Type of randomisation; details of any restriction (such as blocking and block size)	4
Allocation concealment mechanism‡	9	Mechanism used to implement the random allocation sequence§ (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	4
Implementation‡	10	Who generated the random allocation sequence,§ who enrolled participants, and who assigned participants to the sequence of interventions	4
Blinding‡	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	4-5
Similarity of interventions‡	11b	If relevant, description of the similarity of interventions	5-6
Statistical method‡	12a	Statistical methods used to compare groups for primary and secondary outcomes which are appropriate for crossover design (that is, based on within participant comparison)	6-7
Additional analyses‡	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	7
Results:			
Participant flow (a diagram is strongly recommended)†	13a	The numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome, separately for each sequence and period	8
Losses and exclusions†	13b	No of participants excluded at each stage, with reasons, separately for each sequence and period	8
Recruitment†	14a	Dates defining the periods of recruitment and follow-up	8
Trial end†	14b	Why the trial ended or was stopped	8
Baseline data†	15	A table showing baseline demographic and clinical characteristics by sequence and period	4
Numbers analysed†	16	Number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups	8
Outcomes and estimation†	17a	For each primary and secondary outcome, results including estimated effect size and its precision (such as 95% confidence interval) should be based on within participant comparisons.¶ In addition, results for each intervention in each period are recommended	9-10
Binary outcomes‡	17b	For binary outcomes, presentation of both absolute and relative effect sizes is recommended	
Ancillary analyses‡	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing prespecified from exploratory	10-11
Harm‡	19	Describe all important harms or unintended effects in a way that accounts for the design (for specific guidance, see CONSORT for harms <sup>3,5</sup> )	8
Discussion:			
Limitations†	20	Trial limitations, addressing sources of potential bias, imprecision, and if relevant, multiplicity of analyses. Consider potential carry over effects	14
Generalisability†	21	Generalisability (external validity, applicability) of the trial findings	12-13
Interpretation†	22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	12-14
Other information:			
Registration†	23	Registration number and name of trial registry	5
Protocol†	24	Where the full trial protocol can be accessed, if available	
Funding‡	25	Sources of funding and other support (such as supply of drugs), role of funders	14

CONSORT=Consolidated Standards of Reporting Trials.

\*Note: page numbers are optional depending on journal requirements.

†Modified original CONSORT item.

‡Unmodified CONSORT item.

§Random sequence here refers to a list of random orders, typically generated through a computer program. This should not be confused with the sequence of interventions in a randomised crossover trial, for example receiving intervention A before B for an individual trial participant.

¶A within participant comparison takes into account the correlation between measurements for each participant because they act as their own control, therefore measurements are not independent.

BMJ: first published as 10.1136/bmj.14378 on 31 July 2019. Downloaded from http://www.bmj.com/ on 31 July 2019 by guest. Protected by copyright.



### 3. Eigenständigkeitserklärung:

#### **Eigenständigkeitserklärung**

Ich (de Groot, Quintin) erkläre hiermit, dass ich die vorliegende Arbeit selbständig, ohne Mithilfe Dritter und unter Benützung der angegebenen Quellen verfasst habe.

Ort, Datum: Winterthur, 17.05.2020

Unterschrift:



#### 4. Kritische Würdigung der eigenen Arbeit:

**Studie:** Comparison of two commercially available devices to measure nitric oxide lung diffusing capacity in healthy, non-smoking adults.

**Eigene Leistung:**

Meine eigene Leistung an dieser Forschungsarbeit hat schon im ersten Transferpraktikum begonnen. Ich stieg in ein schon geplantes Projekt zwischen den Standorten der Spitäler in Paris, Aachen und Zürich ein. Bei diesem Projekt, welches später als Pilotstudie eine Basis für diese Masterarbeit legte, wurde ein Vergleich zweier DLNO Messgeräte an den oben genannten Standorten durchgeführt. Meine Rolle bei diesem Projekt war die eines Probanden. Auf Basis der Erkenntnisse dieser Pilotstudie wurde binnen des Instituts für Epidemiologie, Biostatistik und Prävention die Forschungsfrage dieser Studie entwickelt. Da ich schon bei der zuvor durchgeführten Pilotstudie teilnahm und daher in diesem Fachgebiet eingeleitet war, nahm ich dieses Projekt gerne als meine Masterarbeit an. Das Studiendesign stand zu Beginn fest, da es um einen Gerätevergleich handelt. Abgesehen vom Design war es mir jedoch möglich, Vorschläge für das detaillierte Vorgehen einzubringen und meine Meinung zu äußern. So hatte ich zu Beginn den Vorschlag, eine aktivitätsbezogene Intervention wie zum Beispiel 20 Kniebeugen miteinzubauen, um den Zusammenhang mit der Physiotherapie herzustellen. Nachdem vonseiten der ZHAW entschieden wurde, dass eine direkte physiotherapeutische Relevanz nicht mehr obligatorisch sei, konnte auf die endgültige Forschungsfrage fokussiert werden. Folglich konnte ich die Forschungsfrage ausarbeiten und einen detaillierten Ablauf in Zusammenarbeit mit meinem Betreuer planen. Mithilfe einer weiteren Kollegin war es möglich, die Messgeräte die Gegenstand der Studie waren, besser kennenzulernen und selbstorganisierte Probemessungen durchzuführen. Die Probandenrekrutierung außerhalb des Kollegen- und Bekanntenkreises des Institutes und des Universitätsspitals lag in meinem Aufgabenbereich. Auch die Durchführung eines großen Teils der Datenerhebung, sowie die Dateneingabe befanden sich in meiner Verantwortung. Nach Abschluss der Datenerhebung und der ersten Auswertungen wurde mit der Fehlersuche und der Problemlösung begonnen. Es mussten potenzielle

Quellen für die Entstehung des gemessenen Unterschiedes erörtert und der jeweilige mögliche Einfluss analysiert werden. Zusammenfassend kann gesagt werden, dass die Rahmenbedingungen sowie die Projektidee von meinem Betreuer gestellt wurden, ich jedoch binnen dieses Rahmens frei war, Ideen und Alternativen mit einzubringen und mitverantwortlich für die Durchführung, Analyse und Interpretation der Studie war.

### **Offene Fragen:**

Die durchgeführte Studie zeigt auf, dass zwei Geräte binnen des Messverfahrens DLNO bei gleichen Individuen und Umgebungsfaktoren deutlich unterschiedliche Ergebnisse erzeugen. Nun stellt sich die Frage, ob zwei identische Geräte an verschiedenen Standorten vergleichbar messen. Falls dies der Fall sein sollte, könnte man Daten poolen, die auf einem identischen Gerät erhoben wurden. Da es momentan nur zwei auf dem Markt frei erhältliche Messgeräte gibt, könnte es trotz des Ergebnisses aus dieser Studie eine Möglichkeit geben, größere Datensätze als Referenzwerte zu generieren. Aufgrund verschiedener Geräte- und Softwareversionen der zwei Hersteller, wäre auch diesbezüglich ein Vergleich sinnvoll.

### **Kritische Betrachtung:**

Offen bleibt zudem die Frage, ob innerhalb dieser Studie eine Einstellung an einem Gerät übersehen wurde, oder ob ein anderer unbekannter Fehler gemacht wurde. Trotz der Überprüfungen der Techniker und mehrfacher Kontrolle, sowie protokollierte Vorgehensweisen und Biomonitoring bleibt immer die Möglichkeit bestehen, dass eine mögliche Fehlerquelle nicht erkannt wurde. Diese Gefahr besteht jedoch in jeder durchgeführten Studie und nur eine Replikationsstudie könnte diese Art potenzieller Fehlerquellen aufdecken oder eliminieren.

Die Durchführung dieser Studie fand im Rahmen des Masterstudiums Physiotherapie der ZHAW statt. Daher war auch der zeitliche Ablauf sehr eng an die Rahmenbedingungen der ZHAW und der nebenbei anfallenden Studienlast gebunden. Prüfungsphasen, Semesterferien und Module, die auf spezifische Abschnitte der Masterarbeit zugeschnitten waren, bestimmten den zeitlichen Ablauf zu einem gewissen Maß. Hätte diese Bindung nicht existiert, wäre es möglich gewesen, diese

Forschungsarbeit in einem kürzeren zeitlichen Rahmen durchzuführen. Ein weiterer Grund, der für eine zwischenzeitliche Zwangspause gesorgt hatte, war eine leere Gasflasche und die entsprechende Lieferzeit der besagten Flasche. Nach einer fünfwöchigen Lieferzeit wurde diese kurz vor Weihnachten 2019 geliefert. Dies führte dazu, dass die Datenerhebung erst Mitte Januar anstelle von November/Dezember beendet werden konnte. Ab Februar entstanden erneut Schwierigkeiten bezüglich Besprechungsterminen und Terminen im Labor, aufgrund der Corona Pandemie.

### **Alternative Vorgehensweise:**

In einer zukünftigen Replikationsstudie könnten verschiedene Maßnahmen getroffen werden, um einen Gerätevergleich zu optimieren. Zum einen wäre es wichtig, bei beiden Geräten die identische Gas-Mischung zu verwenden. Beide Geräte wurden jedoch in dieser Studie auf die jeweilige Gaskonzentration kalibriert, jedoch würde dies ein methodologisch optimierten Vergleich gewährleisten. Des Weiteren sollte eine externe Messmethode verwendet werden, um die Atem-Anhalte-Zeit (BHT) zu regulieren. Dies könnte beispielsweise mithilfe eines akustischen Signals gehandhabt werden. Ein solches Vorgehen würde den Einfluss des Programm-Interfaces minimieren. Die verwendeten Gasflaschen sollten regelmäßig kontrolliert werden und frühzeitig benötigte Gase bestellt werden. Eine weitere mögliche Verbesserung wäre, die Beschränkung der Verantwortlichkeit auf eine Person für die Durchführung der Messung zu kürzen. Dies würde einen potenziellen Einfluss von verschiedenen Messpersonen auf die Datenerhebung beseitigen. Im Rahmen dieser Masterstudie war dies aus Zeitgründen nicht möglich.

### **Relevanz der Masterarbeit:**

Diese Studie hat nur bedingt eine Relevanz für die Physiotherapie. Innerhalb der Pneumologie und der Lungenfunktionsdiagnostik zeigt diese Studie jedoch eine sehr viel größere Relevanz. Wie in der Einleitung der Forschungsarbeit schon beschrieben, könnte dieses Verfahren helfen die Lungendiffusion und Krankheiten, bei denen dieser Prozess eingeschränkt ist, besser zu verstehen. Referenzwerte sind ein essenzieller Bestandteil, um individuelle Ergebnisse klinisch interpretieren zu können. In den

vergangenen Jahren wurden vermehrt Studien zur Generierung von Referenzwerten durchgeführt. Auch wurden diese Ergebnisse schon zusammengeführt, um die Aussagekraft zu vergrößern. Unsere Studie zeigt jedoch, dass dieser Schritt möglicherweise noch nicht gemacht hätte werden sollen. Wir haben aufgedeckt, dass man die Daten von den untersuchten Messgeräten nicht zusammenführen kann. Dies könnte später bei klinischer Anwendung im Ernstfall zu einer verfälschten medizinischen Diagnostik führen.

Die Relevanz für den Fachbereich der Physiotherapie besteht darin, dass dieses Verfahren gerade bei kardio-pulmonalen Patienten häufig einen elementaren Bestandteil darstellt. Daher hilft es zu gewährleisten, dass Patienten beispielsweise mit einer COPD, ein evidenzbasiertes diagnostisches Verfahren erhalten.

Generell ist nach momentanem Forschungsstand der Lungendiffusionsmessung mittels Stickstoffmonoxid noch nicht klar, welche Vorteile diese Messmethode mit sich bringt und ob diese jemals klinisch in Kombination zur DLCO Anwendung findet. Laut aktuellem Wissenstand könnte diese Methode, potenziell zu einem besseren Verständnis des Gasaustausches binnen der Lungenfunktionsdiagnostik führen. Diese Forschungsarbeit zeigt neue Erkenntnisse auf und beantwortet eine wichtige, bisher ungeklärte Forschungsfrage. Insgesamt hilft die durchgeführte Studie das Messverfahren mittels Stickstoffmonoxid besser zu verstehen und bringt damit den potenziellen Mehrwert einen Schritt näher an den Patienten.

### **Persönlicher Bezug:**

Der Fachbereich Lungendiffusion war auch für mich als muskuloskelettaler Physiotherapeut etwas Unbekanntes. Nach langer Einarbeit binnen des ersten Transfermoduls und der beiden Messungen in Paris und Aachen, hat dieses Thema mein Interesse geweckt und ich habe es als sehr abwechslungsreich und erkenntniserweiternd empfunden, binnen der Pneumologie eine Forschungsarbeit durchgeführt zu haben. Meine bisherige Arbeitserfahrung hat sich sehr auf die physiotherapeutische Arbeit in Einzelpraxen konzentriert. Daher stellt diese Masterarbeit mit dem Einblick in den Fachbereich der inneren Medizin einen interdisziplinären, spannenden und herausfordernden Abschnitt des Masterstudiums dar.

## 5. Ethikkommission-Bewilligung:



Universität Zürich  
Epidemiology, Biostatistics and  
Prevention Institute  
Dr. Thomas Radtke  
HRS F 108001  
Hirschengraben 84  
8001 Zürich

Kanton Zürich  
**Kantonale Ethikkommission**



**Prof. Dr. med. Peter Meier-Abt**  
Präsident

**Dr. iur. / lic. phil. Lea Schläpfer**  
Juristin  
Stampfenbachstrasse 121  
Postfach  
8090 Zürich  
Telefon +41 43 259 79 70  
Fax +41 43 259 79 72  
[www.kek.zh.ch](http://www.kek.zh.ch)

19<sup>th</sup> March 2019 / jam

**BASEC -Nr. Req-2019-00226**

**Clarification of responsibility**

**Title: Comparison of two commercially available devices to measure nitric oxide lung diffusing capacity in healthy, non-smoking adults.**

Dear Doctor Radtke

We refer to your submission dated 12/03/2019.

Your research project does not fall within the scope of the Human Research Act (HRA). Therefore, an authorization from the ethics committee is not required.

Kindly note that an invoice in the amount of CHF 200.- will be issued by the cantonal accounts department.

Sincerely yours,

Lea Schläpfer

## 6. Bestätigung der Betreuung:

### **Bestätigung der Betreuungsperson**

Die betreuende Person, (Thomas Radtke) gibt ihr Einverständnis, dass die vorliegende Version als Masterarbeit eingereicht wird.

Ort, Datum: Zürich, 04.05.2020

Unterschrift: 

## 7. Curriculum Vitae des Verfassers:

# Quintin Olivier de Groot

08.11.1993  
Kempten (Allgäu)

Schlosshofstrasse 17 - 8400 Winterthur (Switzerland)  
 Quintindegroot93@gmail.com  
 +31 (0)640328577  
 German/Dutch



## Education

<b>2017 – present</b>	<b>Master of Science in Physiotherapy</b> <i>ZHAW Zurich University of Applied Sciences (Switzerland)</i>
<b>2013 – 2017</b>	<b>Bachelor of Science in Physiotherapy</b> <i>Hogeschool Arnhem en Nijmegen (Netherlands)</i>
<b>2012 – 2013</b>	<b>Work &amp; Travel</b> <i>New Zealand and Southeast-Asia</i>
<b>2010 – 2012</b>	<b>Specialized A-Levels (Social Services)</b> <i>Fachoberschule Kempten (Allgäu) (Germany)</i>
<b>2005 – 2010</b>	<b>Mittlere Reife</b> <i>Staatliche Realschule Kempten (Allgäu) (Germany)</i>

## Work Experience

<b>2020 – present</b>	<b>Physiotherapist at Kantonspital Winterthur</b> <i>Winterthur (Switzerland)</i>
<b>2017 – 2020</b>	<b>Physiotherapist at Medbase Group</b> <i>Uzwil (Switzerland)</i>
<b>Winter 2016/17</b>	<b>Trainee Physiotherapist</b> <i>Wijkgezondheidscentrum Lindenholt (Nijmegen)</i>
<b>Summer 2016</b>	<b>Technical Representative BeNeLux</b> <i>Mammut Deutschland (BeNeLux)</i>
<b>Winter 2015</b>	<b>Trainee Physiotherapist</b> <i>Krankengymnastik Scholten Sulzberg (Germany)</i>

## Courses

<b>present</b>	OMT- Osteopathic manual therapy
<b>2016</b>	Medical Taping
<b>2014</b>	Classical Taping
<b>2013</b>	Reanimation
<b>2010</b>	Extended First-Aid

## Languages

<b>Dutch:</b>	Fluent
<b>German:</b>	Fluent
<b>English:</b>	Advanced
<b>French:</b>	Beginner